www.antibodiesinc.com
orders@antibodiesinc.com
530-758-4400

PRODUCT DATASHEET

Anti-Phospho-Ubiquitin Ser65 Recombinant Chicken Chimeric mAb FL550 Conjugate

Product Details

Available Variants	200 µL (SKU:78-606-FL550)
Conjugate	550
Isotype	IgY
Clone	41H1K3
Gene Name	UBC, UBB
Host Species	Chicken
Concentration	Lot dependent: provided at 0.3-0.5 mg/mL
Format	Purified by affinity chromatography
Physical State	Liquid
Buffer	1X PBS, 0.05% Sodium Azide pH 7.4
Production Notes	This recombinant antibody is a chimeric antibody created by replacing the mouse heavy and light constant regions of clone 41H1K3 with chicken IgY heavy and light constant regions. As such this antibody retains the same binding performance as the original clone 41H1K3 but can be detected using standard anti-chicken secondary antibodies allowing flexibility for multiplexing applications. This antibody is expressed recombinantly in mammalian cells and then affinity purified from the cell culture media and conjugated.
Applications	IHC
Dilution Ranges	IHC: 1:200-1:1000
Species Reactivity	Human, Mouse, Rat
Immunogen	The exact immunogen used to generate this antibody is proprietary information
Specificity	No cross-reactivity reported

Molecular Weight	Multiple bands
Quality Control	Each new lot of antibody is quality control tested by Western blot on mock- and CCCP-treated HeLa lysates and confirmed to stain multiple bands only in CCCP treated lysates
Storage	Aliquot and store at ≤ -20°C for long term storage. For short term storage, store at 2-8°C. For maximum recovery of product, centrifuge the vial prior to removing the cap.
UniProt ID	P0CG48 UniProt Summary: Exists either covalently attached to another protein, or free (unanchored). When covalently bound, it is conjugated to target proteins via an isopeptide bond either as a monomer (monoubiquitin), a polymer linked via different Lys residues of the ubiquitin (polyubiquitin chains) or a linear polymer linked via the initiator Met of the ubiquitin (linear polyubiquitin chains). Polyubiquitin chains, when attached to a target protein, have different functions depending on the Lys residue of the ubiquitin that is linked: Lys-6-linked may be involved in DNA repair; Lys-11-linked is involved in ERAD (endoplasmic reticulum-associated degradation) and in cell-cycle regulation; Lys-29-linked is involved in proteotoxic stress response and cell cycle; Lys-33-linked is involved in kinase modification; Lys-48-linked is involved in protein degradation via the proteasome; Lys-63-linked is involved in endocytosis, DNA-damage responses as well as in signaling processes leading to activation of the transcription factor NF-kappa-B. Linear polymer chains formed via attachment by the initiator Met lead to cell signaling. Ubiquitin is usually conjugated to Lys residues of target proteins, however, in rare cases, conjugation to Cys or Ser residues has been observed. When polyubiquitin is free (unanchored-polyubiquitin), it also has distinct roles, such as in activation of protein kinases, and in signaling. During ubiquitination, the acceptor ubiquitin is positioned in the active site via direct interaction with the E2 ubiquitin-conjugating enzymes such as UBE2R2. As a monoubiquitin, its C-terminal glycine is recognized as a C-degron by Cul2-RING (CRL2) E3 ubiquitin-protein ligase complexes (PubMed:39548056)
Country of Origin	United States
Shipping	Shipped on ice packs
Expiration	24 months from date of receipt
Usage Statement	These antibodies are to be used as research laboratory reagents and are not for use as diagnostic or therapeutic reagents in humans.
NCBI Gene ID	7316

Product Images



Anti-Phospho-Ubiquitin Ser65 Recombinant Chicken Chimeric mAb FL550 Conjugate