www.antibodiesinc.com
orders@antibodiesinc.com
530-758-4400

PRODUCT DATASHEET

Anti-Neuregulin-CRD (Cysteine-Rich Domain, Type III) Antibody FL550 Conjugate

Product Details

Available Variants	200 µL (SKU:75-226-FL550)
Conjugate	FL550 Ex: 550 nm, Em: 575 nm
Isotype	IgG2b
Clone	N126B/31
Gene Name	NRG1
Host Species	Mouse
Concentration	Lot dependent: provided at 0.3-0.5 mg/mL
Format	Purified by Protein A chromatography
Physical State	Liquid
Buffer	PBS with 0.09% azide
Production Notes	Produced by in vitro bioreactor culture of hybridoma line followed by Protein A affinity chromatography and conjugation of purified mAb. Purified mAbs are >90% specific antibody.
Applications	ICC
Species Reactivity	Human, Rat
Immunogen	Fusion protein amino acids 1-75 (membrane-associated N-terminus) of human Neuregulin-1 (accession number Q96IB3) produced recombinantly in E. Coli
Specificity	No cross-reactivity reported
Molecular Weight	30 kDa
Quality Control	Each new lot of antibody is quality control tested on cells overexpressing target protein and confirmed to give the expected staining pattern.

Storage	Aliquot and store at ≤ -20°C for long term storage. For short term storage, store at 2-8°C. For maximum recovery of product, centrifuge the vial prior to removing the cap.
Antibody Registry ID	AB_2939825
UniProt ID	Q96IB3 UniProt Summary: Direct ligand for ERBB3 and ERBB4 tyrosine kinase receptors. Concomitantly recruits ERBB1 and ERBB2 coreceptors, resulting in ligand-stimulated tyrosine phosphorylation and activation of the ERBB receptors. The multiple isoforms perform diverse functions such as inducing growth and differentiation of epithelial, glial, neuronal, and skeletal muscle cells; inducing expression of acetylcholine receptor in synaptic vesicles during the formation of the neuromuscular junction; stimulating lobuloalveolar budding and milk production in the mammary gland and inducing differentiation of mammary tumor cells; stimulating Schwann cell proliferation; implication in the development of the myocardium such as trabeculation of the developing heart. Isoform 10 may play a role in motor and sensory neuron development. Binds to ERBB4. Binds to ERBB3. Acts as a ligand for integrins and binds (via EGF domain) to integrins ITGAV:ITGB3 or ITGA6:ITGB4. Its binding to integrins and subsequent ternary complex formation with integrins and ERRB3 are essential for NRG1-ERBB signaling. Induces the phosphorylation and activation of MAPK3/ERK1, MAPK1/ERK2 and AKT1. Ligand-dependent ERBB4 endocytosis is essential for the NRG1-mediated activation of these kinases in neurons.
Country of Origin	United States
Shipping	Shipped on ice packs
Expiration	12 months from date of receipt
Usage Statement	These antibodies are to be used as research laboratory reagents and are not for use as diagnostic or therapeutic reagents in humans.
NCBI Gene ID	3084

Product Images



Anti-Neuregulin-CRD (Cysteine-Rich Domain, Type III) Antibody FL550 Conjugate