www.antibodiesinc.com
orders@antibodiesinc.com
530-758-4400

PRODUCT DATASHEET

Anti-Kv1.4 Potassium Channel (Extracellular) Antibody FL490 Conjugate

NeuroMab

Product Details

Available Variants	200 µL (SKU:75-316-FL490)
Conjugate	FL490 Ex: 491 nm, Em: 515 nm
Isotype	IgG1
Clone	L71/5
Gene Name	Kcna4
Host Species	Mouse
Concentration	Lot dependent: provided at 0.3-0.5 mg/mL
Format	Purified by Protein A chromatography
Physical State	Liquid
Buffer	PBS with 0.09% azide
Production Notes	Produced by in vitro bioreactor culture of hybridoma line followed by Protein A affinity chromatography and conjugation of purified mAb. Purified mAbs are >90% specific antibody.
Applications	ICC
Species Reactivity	Mouse, Rat
Immunogen	Fusion protein amino acids 336-370 (DDRDLIMALSAGGHSRLLNDTSAPHLENSGHTIFN, extracellular S1-S2 linker) of rat Kv1.4 (accession number P15385) produced recombinantly in E. Coli
Specificity	No cross-reactivity reported
Molecular Weight	98 kDa

Quality Control	Each new lot of antibody is quality control tested on cells overexpressing target protein and confirmed to give the expected staining pattern.
Storage	Aliquot and store at ≤ -20°C for long term storage. For short term storage, store at 2-8°C. For maximum recovery of product, centrifuge the vial prior to removing the cap.
Antibody Registry ID	AB_2940104
UniProt ID	P15385 UniProt Summary: Voltage-gated potassium channel that mediates transmembrane potassium transport in excitable membranes. Forms tetrameric potassium-selective channels through which potassium ions pass in accordance with their electrochemical gradient. The channel alternates between opened and closed conformations in response to the voltage difference across the membrane. Can form functional homotetrameric channels and heterotetrameric channels that contain variable proportions of KCNA1, KCNA2, KCNA4, KCNA5, and possibly other family members as well; channel properties depend on the type of alpha subunits that are part of the channel. Channel properties are modulated by cytoplasmic beta subunits that regulate the subcellular location of the alpha subunits and promote rapid inactivation. In vivo, membranes probably contain a mixture of heteromeric potassium channel complexes, making it difficult to assign currents observed in intact tissues to any particular potassium channel family member. Homotetrameric KCNA4 forms a potassium channel that opens in response to membrane depolarization, followed by rapid spontaneous channel closure. Likewise, a heterotetrameric channel formed by KCNA1 and KCNA4 shows rapid inactivation.
Country of Origin	United States
Shipping	Shipped on ice packs
Expiration	12 months from date of receipt
Usage Statement	These antibodies are to be used as research laboratory reagents and are not for use as diagnostic or therapeutic reagents in humans.
NCBI Gene ID	25469

Product Images



Anti-Kv1.4 Potassium Channel (Extracellular) Antibody FL490 Conjugate