www.antibodiesinc.com
orders@antibodiesinc.com
530-758-4400

PRODUCT DATASHEET

Anti-Homer1 Antibody FL650 Conjugate

Product Details

Available Variants	200 µL (SKU:75-454-FL650)
Conjugate	FL650 Ex: 655 nm, Em: 676 nm
Isotype	IgG1
Clone	L113/130
Gene Name	Homer1 Vesl1
Host Species	Mouse
Concentration	Lot dependent: provided at 0.3-0.5 mg/mL
Format	Purified by Protein A chromatography
Physical State	Liquid
Buffer	PBS with 0.09% azide
Production Notes	Produced by in vitro bioreactor culture of hybridoma line followed by Protein A affinity chromatography and conjugation of purified mAb. Purified mAbs are >90% specific antibody.
Applications	ICC, IHC
Species Reactivity	Human, Mouse, Rat
Immunogen	Fusion protein amino acids 121-363 (C-terminus) of mouse Homer1 (accession number Q9Z2Y3) produced recombinantly in E. Coli
Specificity	No cross-reactivity reported
Molecular Weight	50 kDa
Quality Control	Each new lot of antibody is quality control tested by western blot on rat whole brain lysate and confirmed to stain the expected molecular weight band.
Storage	Aliquot and store at ≤ -20°C for long term storage. For short term storage, store at 2-8°C. For maximum recovery of product, centrifuge the vial prior to

	removing the cap.
Antibody Registry ID	AB_2940511
UniProt ID	Q9Z2Y3 UniProt Summary: Postsynaptic density scaffolding protein. Binds and cross-links cytoplasmic regions of GRM1, GRM5, ITPR1, DNM3, RYR1, RYR2, SHANK1 and SHANK3. By physically linking GRM1 and GRM5 with ER-associated ITPR1 receptors, it aids the coupling of surface receptors to intracellular calcium release. May also couple GRM1 to PI3 kinase through its interaction with AGAP2. Isoform 1 regulates the trafficking and surface expression of GRM5. Differentially regulates the functions of the calcium activated channel ryanodine receptors RYR1 and RYR2. Isoform 1 decreases the activity of RYR2, and increases the activity of RYR1, whereas isoform 5 counteracts the effects by competing for binding sites. Isoform 3 regulates the trafficking and surface expression of GRM5. Isoform 5 acts as a natural dominant negative, in dynamic competition with constitutively expressed isoform 1, isoform 2 and isoform 3 to regulate synaptic metabotropic glutamate function. Isoform 5, may be involved in the structural changes that occur at synapses during long-lasting neuronal plasticity and development. Forms a high-order complex with SHANK1, which in turn is necessary for the structural and functional integrity of dendritic spines. Negatively regulates T cell activation by inhibiting the calcineurin-NFAT pathway. Acts by competing with calcineurin/PPP3CA for NFAT protein binding, hence preventing NFAT activation by PPP3CA.
Country of Origin	United States
Shipping	Shipped on ice packs
Expiration	12 months from date of receipt
Usage Statement	These antibodies are to be used as research laboratory reagents and are not for use as diagnostic or therapeutic reagents in humans.
NCBI Gene ID	26556

Product Images



Anti-Homer1 Antibody FL650 Conjugate