

Anti-NMDA NR2B Subunit (Tyr1336) Antibody




Pooled Serum

Product Details


Available Variants	100 µL (SKU:p1516-1336) 25 µL (SKU:p1516-1336t)
Conjugate	Unconjugated
Isotype	IgG
Gene Name	GRIN2B
Host Species	Rabbit
Format	Antigen Affinity Purified from Pooled Serum
Physical State	Liquid
Buffer	10 mM HEPES (pH 7.5), 150 mM NaCl, 100 µg per ml BSA and 50% glycerol.
Production Notes	Prepared from pooled rabbit serum by affinity purification via sequential chromatography on phospho and non-phosphopeptide affinity columns.
Applications	ELISA, IHC, WB
Dilution Ranges	WB: 1:1000 WB Brain: 1:1000 IHC: 1:400 ICC: 1:100
Species Reactivity	Mouse, Rat
Immunogen	Synthetic phospho-peptide corresponding to amino acid residues surrounding Tyr1336 of the NR2B subunit of the rat NMDA receptor, conjugated to keyhole limpet hemocyanin (KLH).
Specificity	Specific for endogenous levels of the ~180 kDa NMDAR NR2B-subunit protein phosphorylated at Tyr1336 . Immunolabeling is completely eliminated by treatment with λ-phosphatase.
Molecular Weight	180 kDa

Quality Control	Western blots performed on each lot.
Storage	Storage at -20°C is recommended, as aliquots may be taken without freeze/thawing due to presence of 50% glycerol. Stable for at least 1 year at -20°C.
Antibody Registry ID	AB_2492181
UniProt ID	<u>Q00960</u>
Shipping	Blue Ice
Expiration	After date of receipt, stable for at least 1 year at -20°C.
Usage Statement	For research use only. Not intended for therapeutic or diagnostic use. Use of all products is subject to our terms and conditions, which can be viewed on our website.
Expected Reactivity	Human, Non-Human Primate
Post Translational Modification Type	Phosphorylated
Post Translational Modification Residue	Tyr1336
NCBI Gene ID	24410

Product Images



Western blot of rat hippocampal lysate showing specific immunolabeling of the ~180 kDa NR2B subunit phosphorylated at Tyr1336 in the first lane (-). Phosphospecificity is shown in the second lane (+) where immunolabeling is completely eliminated by blot treatment with lambda phosphatase (λ -Ptase, 1200 units for 30 min).



Immunostaining of mouse dentate gyrus 48 hours post TMT treatment showing NR2B when phosphorylated at Tyr1336 (green, 1:400). The blue is staining nuclei using DAPI. Photo Courtesy of Rob Wine.

Product Page URL: www.antibodiesinc.com/products/anti-nmda-nr2b-subunit-tyr1336-antibody-p1516-1336



Created on 04. September 2026 | All information without guarantee