www.antibodiesinc.com
orders@antibodiesinc.com
530-758-4400

PRODUCT DATASHEET

Anti-c-Cbl (C-terminal region) Antibody


phosphosolutions ANTIBODIES THAT WORK

Product Details

Available Variants	100 µL (SKU:CM1591)
Conjugate	Unconjugated
Isotype	IgG1
Clone	M159
Gene Name	CBL
Host Species	Mouse
Format	Protein A Purified
Physical State	Liquid
Buffer	PBS + 1 mg/ml BSA, 0.05% NaN3 and 50% glycerol
Applications	ELISA, WB
Dilution Ranges	WB: 1:1000
Species Reactivity	Human, Mouse, Rat
Immunogen	Clone (M159) was generated from a synthetic peptide (coupled to KLH) corresponding to amino acid residues in the C-terminal region of human c-Cbl. This sequence is highly conserved in rat and mouse c-Cbl.
Specificity	The antibody detects a 120 kDa* protein in human Jurkat cells.
Molecular Weight	120
Quality Control	Western blots performed on each lot.
Storage	Storage at -20°C is recommended, as aliquots may be taken without freeze/thawing due to presence of 50% glycerol. Stable for at least 1 year at -20°C.
UniProt ID	P22681 UniProt Summary: E3 ubiquitin-protein ligase that acts as a negative regulator

	of many signaling pathways by mediating ubiquitination of cell surface receptors. Accepts ubiquitin from specific E2 ubiquitin-conjugating enzymes, and then transfers it to substrates promoting their degradation by the proteasome. Recognizes activated receptor tyrosine kinases, including KIT, FLT1, FGFR1, FGFR2, PDGFRA, PDGFRB, CSF1R, EPHA8 and KDR and mediates their ubiquitination to terminate signaling. Recognizes membrane-bound HCK, SRC and other kinases of the SRC family and mediates their ubiquitination and degradation. Ubiquitinates EGFR and SPRY2. Involved in LAG3-mediated inhibition of TCR signaling: following ligand-binding to LAG3, catalyzes 'Lys-63'-linked ubiquitination of LAG3, unleashing the LAG3 C-terminus from the membrane, and initiating a signaling that prevents TCR activation. Ubiquitinates NECTIN1 following association between NECTIN1 and herpes simplex virus 1/HHV-1 envelope glycoprotein D, leading to NECTIN1 removal from cell surface. Participates in signal transduction in hematopoietic cells. Plays an important role in the regulation of osteoblast differentiation and apoptosis. Essential for osteoclastic bone resorption. The 'Tyr-731' phosphorylated form induces the activation and recruitment of phosphatidylinositol 3-kinase to the cell membrane in a signaling pathway that is critical for osteoclast function. In association with CBLB, required for proper feedback inhibition of ciliary platelet-derived growth factor receptor-alpha (PDGFRA) signaling pathway via ubiquitination and internalization of PDGFRA.
Country of Origin	United States
Shipping	Blue Ice
Expiration	After date of receipt, stable for at least 1 year at -20°C.
Usage Statement	For research use only. Not intended for therapeutic or diagnostic use. Use of all products is subject to our terms and conditions, which can be viewed on our website.
NCBI Gene ID	867

Product Images



Western blot analysis of human Jurkat cells untreated (lanes 1 & 3) or treated with pervanadate (1 mM) for 30 minutes (lanes 2 & 4). The blot was probed with anti-c-Cbl (CM1591; lanes 1 & 2) or anti-c-Cbl (Tyr-700) (CM1611; lanes 3 & 4).