

Anti-Myosin Light Chain (Ser-1), Phosphospecific Antibody




Product Details


Available Variants	100 µL (SKU:MP3461)
Conjugate	Unconjugated
Isotype	IgG
Gene Name	MYL12A
Host Species	Rabbit
Format	Antigen Affinity Purified
Physical State	Liquid
Buffer	PBS + 1 mg/ml BSA, 0.05% NaN ₃ and 50% glycerol
Applications	ELISA, ICC, WB
Dilution Ranges	WB: 1:1000 ICC: 1:100
Species Reactivity	Bovine, Human, Mouse, Rat
Immunogen	Phospho-MLC (Ser-1) synthetic peptide (coupled to KLH) corresponding to amino acid residues surrounding Ser-1 in bovine myosin regulatory light chain 9 (Myl9). This peptide sequence is highly conserved in human, rat, and mouse Myl9, and in other smooth muscle and non-muscle MLCs.
Specificity	The antibody detects a 20 kDa* band corresponding to the molecular weight of myosin light chain in human A431 treated with EGF or calyculin A. In addition, the antibody detects the 20 kDa myosin light chain in mouse C2C12, rat PC12, and rabbit fibroblasts treated with calyculin A.
Molecular Weight	20
Quality Control	Western blots performed on each lot.

Storage	Storage at -20°C is recommended, as aliquots may be taken without freeze/thawing due to presence of 50% glycerol. Stable for at least 1 year at -20°C.
UniProt ID	<u>P19105</u>
Country of Origin	United States
Shipping	Blue Ice
Expiration	After date of receipt, stable for at least 1 year at -20°C.
Usage Statement	For research use only. Not intended for therapeutic or diagnostic use. Use of all products is subject to our terms and conditions, which can be viewed on our website.
Post Translational Modification Type	Phosphorylated
Post Translational Modification Residue	Ser-1

Product Images



Western blot analysis of C2C12 cells untreated (lanes 1, 3, 5, & 7) or treated with Lambda phosphatase (lanes 2, 4, 6, & 8). The blots were probed with monoclonal anti-MLC20 (clone MY-21) (lanes 1 & 2), polyclonal anti-MLC (Ser-19) phosho-specific (lanes 3 & 4), anti-MLC (Ser-1) phosho-specific (lanes 5 & 6), or anti-MLC (a.a. 11-22) (lanes 7 & 8).



Immunocytochemical labeling of phosphorylated MLC in paraformaldehyde fixed A7r5 cells. The cells were dual-labeled with anti-MLC (MM3441; middle) and anti-MLC (MP4201; top left), anti-MLC (Ser-19) (MP4221; middle left) and anti-MLC (Ser-1) (MP3461; bottom left). Goat anti-Mouse DyLight® 488 and Goat anti-Rabbit DyLight® 594 were used for detection of primary antibodies. The overlay of staining patterns are shown to the right.

Product Page URL: www.antibodiesinc.com/products/anti-myosin-light-chain-ser-1-phosphospecific-antibody-mp3461



Created on 10. September 2026 | All information without guarantee