www.antibodiesinc.com
orders@antibodiesinc.com
530-758-4400

PRODUCT DATASHEET

Anti-Tafazzin Antibody (BSA and Azide free)

Product Details

Available Variants	100 µL (SKU:75-197-CF)
Conjugate	Unconjugated
Isotype	IgG2b
Clone	N173B/13
Gene Name	TAZ EFE2 G4.5
Host Species	Mouse
Concentration	Lot dependent: provided at 0.6-1.0 mg/mL
Format	Purified by Protein A chromatography
Physical State	Liquid
Buffer	PBS
Production Notes	Produced by in vitro bioreactor culture of hybridoma line followed by Protein A affinity chromatography and buffer exchanged into 1X PBS. Purified mAbs are >90% specific antibody.
Applications	ELISA, ICC, WB
Species Reactivity	Human
Immunogen	Fusion protein amino acids 1-262 (full-length) of human Tafazzin with exon 5 deleted (accession number Q16635-3) produced recombinantly in E. Coli
Specificity	No cross-reactivity reported
Molecular Weight	30 kDa
Quality Control	Each new lot of antibody is quality control tested on cells overexpressing target protein and confirmed to give the expected staining pattern.
Storage	Aliquot and store at ≤ -20°C for long term storage. For short term storage, store at 2-8°C. For maximum recovery of product, centrifuge the vial prior to

	removing the cap.
UniProt ID	Q16635 UniProt Summary: Acyltransferase required to remodel newly synthesized phospholipid cardiolipin (1',3'-bis-[1,2-diacyl-sn-glycero-3-phospho]-glycerol or CL), a key component of the mitochondrial inner membrane, with tissue specific acyl chains necessary for adequate mitochondrial function. Its role in cellular physiology is to improve mitochondrial performance. CL is critical for the coassembly of lipids and proteins in mitochondrial membranes, for instance, remodeling of the acyl groups of CL in the mitochondrial inner membrane affects the assembly and stability of respiratory chain complex IV and its supercomplex forms. Catalyzes the transacylation between phospholipids and lysophospholipids, with the highest rate being between phosphatidylcholine (1,2-diacyl-sn-glycero-3-phosphocholine or PC) and CL. Catalyzes both 1-acyl-sn-glycero-3-phosphocholine (lysophosphatidylcholine or LPC) reacylation and PC-CL transacylation, that means, it exchanges acyl groups between CL and PC by a combination of forward and reverse transacylations. Also catalyzes transacylations between other phospholipids such as phosphatidylethanolamine (1,2-diacyl-sn-glycero-3-phosphoethanolamine or PE) and CL, between PC and PE, and between PC and phosphatidate (1,2-diacyl-sn-glycero-3-phosphate or PA), although at lower rate. Not regiospecific, it transfers acyl groups into any of the sn-1 and sn-2 positions of the monolysocardiolipin (MLCL), which is an important prerequisite for uniformity and symmetry in CL acyl distribution. Cannot transacylate dilysocardiolipin (DLCL), thus, the role of MLCL is limited to that of an acyl acceptor. CoA-independent, it can reshuffle molecular species within a single phospholipid class. Redistributes fatty acids between MLCL, CL, and other lipids, which prolongs the half-life of CL. Its action is completely reversible, which allows for cyclic changes, such as fission and fusion or bending and flattening of the membrane. Hence, by contributing to the flexibility of the lipid composition, it plays an important role in the dynamics of mitochondria membranes. Essential for the final stage of spermatogenesis, spermatid individualization. Required for the initiation of mitophagy. Required to ensure progression of spermatocytes through meiosis. Exon 7 of human tafazzin is essential for catalysis.
Country of Origin	United States
Shipping	Shipped on ice packs
Expiration	12 months from date of receipt
Usage Statement	These antibodies are to be used as research laboratory reagents and are not for use as diagnostic or therapeutic reagents in humans.
NCBI Gene ID	6901

Product Images



Anti-Tafazzin Antibody (BSA and Azide free)

Product Page URL: www.antibodiesinc.com/products/anti-tafazzin-antibody-n173b-13-75-197-cf

Created on 21. November 2026 | All information without guarantee