www.antibodiesinc.com
orders@antibodiesinc.com
530-758-4400

PRODUCT DATASHEET

Anti-Flrt3 Antibody (BSA and Azide free)

Product Details

Available Variants	100 µL (SKU:75-328-CF)
Conjugate	Unconjugated
Isotype	IgG1
Clone	N353/13
Gene Name	Flrt3 Kiaa1469
Host Species	Mouse
Concentration	Lot dependent: provided at 0.6-1.0 mg/mL
Format	Purified by Protein A chromatography
Physical State	Liquid
Buffer	PBS
Production Notes	Produced by in vitro bioreactor culture of hybridoma line followed by Protein A affinity chromatography and buffer exchanged into 1X PBS. Purified mAbs are >90% specific antibody.
Applications	ELISA, ICC
Species Reactivity	Mouse
Immunogen	Fusion protein amino acids 29-526 (extracellular domain) of mouse Flrt3 (accession number Q8BGT1) produced recombinantly in E. Coli
Specificity	No cross-reactivity against Flrt1 or Flrt2
Molecular Weight	73 kDa
Quality Control	Each new lot of antibody is quality control tested on cells overexpressing target protein and confirmed to give the expected staining pattern.
Storage	Aliquot and store at ≤ -20°C for long term storage. For short term storage, store at 2-8°C. For maximum recovery of product, centrifuge the vial prior to

	removing the cap.
UniProt ID	Q8BGT1 UniProt Summary: Functions in cell-cell adhesion, cell migration and axon guidance, exerting an attractive or repulsive role depending on its interaction partners. Plays a role in the spatial organization of brain neurons. Plays a role in vascular development in the retina. Plays a role in cell-cell adhesion via its interaction with ADGRL3 and probably also other latrophilins that are expressed at the surface of adjacent cells. Interaction with the intracellular domain of ROBO1 mediates axon attraction towards cells expressing NTN1. Mediates axon growth cone collapse and plays a repulsive role in neuron guidance via its interaction with UNC5B, and possibly also other UNC-5 family members. Promotes neurite outgrowth (in vitro). Mediates cell-cell contacts that promote an increase both in neurite number and in neurite length. Plays a role in the regulation of the density of glutamaergic synapses. Plays a role in fibroblast growth factor-mediated signaling cascades. Required for normal morphogenesis during embryonic development, but not for normal embryonic patterning. Required for normal ventral closure, headfold fusion and definitive endoderm migration during embryonic development. Required for the formation of a normal basement membrane and the maintenance of a normal anterior visceral endoderm during embryonic development.
Country of Origin	United States
Shipping	Shipped on ice packs
Expiration	12 months from date of receipt
Usage Statement	These antibodies are to be used as research laboratory reagents and are not for use as diagnostic or therapeutic reagents in humans.
NCBI Gene ID	71436

Product Images



Anti-Flrt3 Antibody (BSA and Azide free)

Created on 08. November 2026 | All information without guarantee