

Anti-Ubiquilin 2 Antibody




Product Details

Available Variants	100 µg (SKU:M-1656-100)
Conjugate	Unconjugated
Isotype	IgG1
Clone	6H9
Host Species	Mouse
Format	IgG
Physical State	Lyophilized
Buffer	Lyophilized from PBS buffer pH 7.2-7.6 with 0.1% trehalose, and sodium azide
Production Notes	Protein G purified
Applications	ICC, WB
Dilution Ranges	WB: 1:1000-1:2000 ICC: 1:500-1:1000
Species Reactivity	Human, Mouse
Immunogen	Recombinant human ubiquilin 2 expressed and purified from E. coli.
Specificity	In primary mouse neuron and glia cell culture, endogenous ubiquilin 2 appears as a weak band at 68 kDa in all transduced and non-transduced cells, indicating low endogenous expression of mouse ubiquilin 2. Strong bands are seen in cells transduced with human wild type or mutant ubiquilin 2. Small proteins which run at 50 kDa in these cells are the fragments of ubiquilin 2. Note, ubiquilin 2 runs at ~66 kDa in human Hela cells and 68 kDa in rodent 3T3 cells. The antibody has also been used successfully for immunocytochemistry.
Molecular Weight	~66 kDa in human Hela cells and 68 kDa in rodent 3T3 cells
Storage	Spin vial briefly before opening. Reconstitute with 100 µL sterile-filtered, ultrapure water to achieve a 1 mg/mL concentration. Centrifuge to remove any


insoluble material. Aliquot and store at -20°C for up to six months after date of receipt. Avoid freeze-thaw cycles.

UniProt ID	<u>Q9UHD9</u>
Country of Origin	United States
Shipping	25°C (ambient)
Expiration	12 months after date of receipt (unopened vial).
Usage Statement	For research use only.


Product Images



Western blot analysis of untransfected primary mouse neuron and glia cell cultures (lane 1), the same cells transduced with human ubiquilin 2 wild type (lane 2), with ubiquilin 2 P506T mutant (lane 3), with ubiquilin 2 P497S mutant (lane 4), with enhanced GFP control (lane 5), in HeLa cells (lane 6) and 3T3 cells (lane 7).



HeLa cell cultures stained with M-1656-100 (green) and our chicken polyclonal antibody to vimentin: C-1409-50 (red). In most individual cells ubiquilin 2 is present diffusely in the cytoplasm of cells, though some cells show enrichment of the protein in spherical autophagosome-like structure.



Left: Analysis of ubiquitin 2 expression in NIH-3T3 cell culture by **Immunocytochemistry**. Cells were stained with mouse antibody to ubiquitin 2 (green, 1:1,000), and co-stained with chicken antibody to lamin A/C (C-1698-100, red, 1:5,000). Blue: DAPI nuclear stain. Cells were treated with 50 μ M of chloroquine, an inhibitor of autophagy, for 16 hours prior to staining. The ubiquitin 2 antibody reveals punctate staining of ubiquitin 2 protein accumulated in lysosomes in the cytoplasm, while the lamin A/C antibody stains the nuclear lamina. **Right: Western blot** analysis of tissue and cell lysates using mouse antibody to ubiquitin 2 (green, 1:1,000). [1] protein standard, [2] NIH-3T3, [3] C6, [4] HEK293, [5] HeLa, [6] SH-SY5Y, [7] rat whole brain, and [8] mouse whole brain. The band at 65-70 kDa corresponds to ubiquitin 2 protein, which is known to differ between human and rodent species.

Product Page URL: www.antibodiesinc.com/products/anti-ubiquitin-2-antibody-m-1656-100



Created on 12. September 2026 | All information without guarantee