


Anti-Sortilin Antibody




Product Details

Available Variants	100 µL (SKU:R-150-100)
Conjugate	Unconjugated
Isotype	Mixed
Host Species	Rabbit
Format	Whole serum
Physical State	Lyophilized
Applications	ELISA, IHC
Dilution Ranges	IHC: 1:1000 ELISA: 1:5000
Species Reactivity	Human, Mouse, Rat
Immunogen	Extracellular domain of glycosylated human sortilin produced in CHO cells was used as the immunogen.
Specificity	Specificity was demonstrated by immunohistochemistry. Reacts with human, rat and mouse Sortilin. Other species not yet tested.
Storage	Spin vial briefly before opening. Reconstitute in 100 µL sterile-filtered, ultrapure water. Centrifuge to remove any insoluble material. After reconstitution keep aliquots at -20°C for a higher stability, and at 2-8°C with an appropriate antibacterial agent. Glycerol (1:1) may be added for an additional stability. Avoid repetitive freeze/thaw cycles.
UniProt ID	Q99523
Country of Origin	United States
Shipping	25°C (ambient)
Expiration	12 months after date of receipt (unopened vial).


Product Images



IHC on rat DRG (free floating cryo section) using Rabbit antibody to Sortilin (neurotensin receptor 3): whole serum (R-150-100) at a dilution of 1: 1000.



IHC on rat spinal cord (free floating cryo section) using Rabbit antibody to Sortilin (neurotensin receptor 3): whole serum (R-150-100) at a dilution of 1: 1000.



IHC on rat trigeminal nerve (free floating cryo section) using Rabbit antibody to Sortilin (neurotensin receptor 3): whole serum (R-150-100) at a dilution of 1: 1000.



Created on 12. September 2026 | All information without guarantee