

Anti-E-Cadherin (Cytoplasmic) (M168) Recombinant Chicken Chimeric mAb




Product Details


Available Variants	100 µL (SKU:78-608) 20 µL (SKU:78-608-020)
Conjugate	Unconjugated
Isotype	IgY
Clone	M168
Gene Name	CDH1
Host Species	Chicken
Concentration	1 mg/mL
Format	Purified by affinity chromatography
Physical State	Liquid
Buffer	1X PBS, 0.05% Sodium Azide pH 7.4
Applications	ICC, IHC, IP, WB
Dilution Ranges	WB: 1:1000-1:3000 IHC: 1:250-1:500 ICC: 1:250-1:1000 IP: 1:100
Species Reactivity	Human, Mouse, Rat
Immunogen	Clone (M168) was generated from a mouse recombinant E-Cadherin protein containing amino acids in the cytoplasmic region. This sequence is highly conserved in human and rat E-cadherin.
Specificity	This E-cadherin antibody detects a 120 kDa E-cadherin protein in human A431 cells, and does not cross-react with VE-cadherin or N-cadherin.

Molecular Weight	120 kDa
Quality Control	Each new lot of antibody is quality control tested by Western blot on MCF-7 cell lysates and confirmed to stain the expected molecular weight band.
Storage	Aliquot and store at $\leq -20^{\circ}\text{C}$ for long term storage. For short term storage, store at $2-8^{\circ}\text{C}$. For maximum recovery of product, centrifuge the vial prior to removing the cap.
UniProt ID	P09803
Country of Origin	United States
Shipping	Blue Ice
Expiration	24 months from date of receipt
Usage Statement	These antibodies are to be used as research laboratory reagents and are not for use as diagnostic or therapeutic reagents in humans.

Product Images



Immunofluorescent staining of MCF-7 and HeLa cells using Aves Labs anti-E-Cadherin (Cytoplasmic) chicken recombinant antibody [M168](Cat No. 78-608). The cells were fixed with 3% paraformaldehyde, washed with PBS, and blocked/permeabilized with 5% non-fat milk powder in PBST. The anti-E-Cadherin antibody (Cat. 78-608) was incubated with the sample at a dilution of 1:1000 (1 ug/mL) at 4C overnight. After washing with PBST, the cells were stained using goat anti-chicken IgY CF488 secondary antibody (Cat No. 80-1001-488) (green). The cells were mounted with Antibodies Incorporated Fluoroshield with DAPI mounting medium (Cat No. AR-6501). DAPI nuclear stain (blue) shows cell nuclei. The staining revealed the characteristic membranous localization of E-Cadherin in MCF-7 cells, with no staining observed in E-Cadherin negative HeLa cells.



Western blot detection of E-Cadherin in MCF-7 (lane 1), HepG2 (lane 2) and HeLa (lane 3) cell lysates using Aves Labs anti-E-Cadherin (Cytoplasmic) chicken recombinant antibody [M168](Cat No. 78-608). 4-20% Tris-Glycine gel was loaded with 20 ug of MCF-7, HepG2 and HeLa cell lysates. Proteins were transferred to PVDF membrane and blocked with 5% non-fat milk powder. Anti-E-Cadherin chicken antibody was incubated with the membrane at a concentration of 1:1000 (1 ug/ml) and then detected via chemiluminescence (30 seconds) using a goat anti-Chicken H+L, HRP conjugated secondary antibody. The anti-E-Cadherin chicken recombinant antibody specifically detected endogenous E-Cadherin in E-Cadherin-positive MCF-7 and HepG2 cell lysates, but not in E-Cadherin-negative HeLa cell lysates, at the expected molecular weight of ~120 kDa. GAPDH was used as a loading control.

Product Page URL: www.antibodiesinc.com/products/anti-e-cadherin-cytoplasmic-recombinant-antibody-m168-78-608



Created on 28. July 2026 | All information without guarantee